



Ballerine® Update:

Global Real-World Experience Q3/20 IUB™ Ballerine® MIDI

101,583 Total Estimated Insertions¹


Event	Rate in %
Expulsion	3.53%
Pregnancy	0.49%
Ectopic Pregnancy	0.01%
Perforation	0.06%

>90% LESS RISK^{3,4}

compared to T-IUDs

40% LESS RISK⁵

compared to T-IUDs




T-IUD:
5%²



T-Cu IUD:
1%²



T-IUD:
>0.26%^{3,4}



T-IUD:
1:909⁵

¹ based on accumulated data from Q1 2015 – Q3 2020 on total estimated insertions and OCON's ongoing vigilance reporting

² Aoun et al. Effects of age, parity, and device type on complications and discontinuation of intrauterine devices. *Obstetrics & Gynecology* (123:3) 2014, 585-592.

³ <https://www.sciencedirect.com/science/article/pii/S0010782415000128>

⁴ [https://www.fertstert.org/article/S0015-0282\(00\)01605-8/pdf](https://www.fertstert.org/article/S0015-0282(00)01605-8/pdf)

⁵ Heinemann et al. Risk of Uterine Perforation with Levonorgestrel-Releasing and Copper Intrauterine Devices in the European. Active Surveillance Study on Intrauterine Devices. *Contraception* (2015), doi: 10.1016/j.contraception.2015.01.007.